



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12P 21/00, C12N 1/20, 1/21, 5/12, 15/00		A1	(11) International Publication Number: WO 99/27128
			(43) International Publication Date: 3 June 1999 (03.06.99)
(21) International Application Number: PCT/US98/25096			(74) Agent: FISH, Andrew, L.; Dechert Price & Rhoads, 4000 Bell Atlantic Tower, 1717 Arch Street, Philadelphia, PA 19103-2793 (US). (81) Designated States: JP, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i>
(22) International Filing Date: 24 November 1998 (24.11.98)			
(30) Priority Data: 08/977,553 25 November 1997 (25.11.97) US			
(71) Applicant: SMITHKLINE BEECHAM CORPORATION [US/US]; One Franklin Plaza, Philadelphia, PA 19103 (US).			
(72) Inventors: PALMER, Leslie, M.; 2820 Eagleville Road, Audubon, PA 19407 (US). FEDON, Jason, C.; Apartment 2, 106 Fairfield Lane, Strafford, PA 19087 (US). WARREN, Richard, L.; 825 Cathcart Road, Blue Bell, PA 19422 (US). TRAINI, Christopher, M.; 50 Potter Court, Media, PA 19063 (US). WANG, Min; 302 Centennial Drive, Blue Bell, PA 19422 (US). JAWORSKI, Deborah, D.; 214 Willowbrook Drive, Norristown, PA 19403 (US). MOONEY, Jeffrey; 143 Winged Foot Court, Limerick, PA 19469 (US). DEBOUCK, Christine; 667 Pugh Road, Wayne, PA 19087 (US). ZHONG, YiYi; Apartment D305, 2828 Egypt Road, Audubon, PA 19403 (US).			
(54) Title: ribH			
(57) Abstract The invention provides ribH polypeptides and polynucleotides encoding ribH polypeptides and methods for producing such polypeptides by recombinant techniques. Also provided are methods for utilizing ribH polypeptides to screen for antibacterial compounds.			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

ribH

FIELD OF THE INVENTION

5 This invention relates to newly identified polynucleotides and polypeptides, and their production and uses, as well as their variants, agonists and antagonists, and their uses. In particular, the invention relates to novel polynucleotides and polypeptides of the lumazine synthase (β -subunit) family, hereinafter referred to as "ribH".

BACKGROUND OF THE INVENTION

10 It is particularly preferred to employ Staphylococcal genes and gene products as targets for the development of antibiotics. The Staphylococci make up a medically important genera of microbes. They are known to produce two types of disease, invasive and toxigenic. Invasive
15 infections are characterized generally by abscess formation effecting both skin surfaces and deep tissues. *S. aureus* is the second leading cause of bacteremia in cancer patients. Osteomyelitis, septic arthritis, septic thrombophlebitis and acute bacterial endocarditis are also relatively common. There are at least three clinical conditions resulting from the toxigenic properties of Staphylococci. The manifestation of these diseases result from the actions of exotoxins as opposed to tissue invasion and
20 bacteremia. These conditions include: Staphylococcal food poisoning, scalded skin syndrome and toxic shock syndrome.

 Riboflavin (vitamin B2) is a member of the B complex of vitamins which function as coenzymes in metabolic reactions. Riboflavin has two coenzyme forms, flavin mononucleotide (FMN) and flavin adenine dinucleotide (FAD) which act in oxidation-reduction reactions such
25 as the cytochrome system of electron transport and the oxidative degradation of pyruvate, fatty acids and amino acids. The first committed step in the biosynthesis of riboflavin is the opening of the imidazole ring of GTP. In the presence of 3 H₂O and Mg⁺⁺, the C-8 of GTP is released as formate accompanied by the release of pyrophosphate by the action of GTP cyclohydrolase II (GCH2; EC 3.5.4.25). This enzyme function is encoded by *ribA* in bacteria and *rib1* in yeast
30 species. Through a series of steps, involving 3,4-dihydroxy-2-butanone 4 phosphate synthase (*ribA*), 6,7-dimethyl-8-ribityllumazine synthase (*ribH*), riboflavin synthase (*ribB*), pyrimidine deaminase and pyrimidine reductase (*ribG*), enzymes encoded by genes within the riboflavin

biosynthesis operon, riboflavin is formed. Because the genes required for riboflavin biosynthesis are present in many pathogenic microorganisms, and since riboflavin biosynthesis has shown to be required for virulence in the swine pathogen *Actinobacillus pleuropneumoniae* (Fuller, TE, *et al.* (1996) A riboflavin auxotroph of *Actinobacillus pleuropneumoniae* is
5 attenuated in swine. Infect. Immun. 64:4659-4664), these gene products represent broad spectrum antibacterial as well as antifungal targets.

The frequency of *Staphylococcus aureus* infections has risen dramatically in the past few decades. This has been attributed to the emergence of multiply antibiotic resistant strains and an increasing population of people with weakened immune systems. It is no longer uncommon to
10 isolate *Staphylococcus aureus* strains which are resistant to some or all of the standard antibiotics. This phenomenon has created a demand for both new anti-microbial agents, vaccines, and diagnostic tests for this organism.

Clearly, there exists a need for factors, such as the ribH embodiments of the invention, that have a present benefit of being useful to screen compounds for antibiotic activity. Such factors are
15 also useful to determine their role in pathogenesis of infection, dysfunction and disease. There is also a need for identification and characterization of such factors and their antagonists and agonists to find ways to prevent, ameliorate or correct such infection, dysfunction and disease.

Certain of the polypeptides of the invention possess amino acid sequence homology to a known *Bacillus amyloliquefaciens* ribH protein.
20

SUMMARY OF THE INVENTION

It is an object of the invention to provide polypeptides that have been identified as novel ribH polypeptides by homology between the amino acid sequence set out in Table 1 [SEQ ID NO: 2]
25 and a known amino acid sequence or sequences of other proteins such as *Bacillus amyloliquefaciens* ribH protein.

It is a further object of the invention to provide polynucleotides that encode ribH polypeptides, particularly polynucleotides that encode the polypeptide herein designated ribH.

In a particularly preferred embodiment of the invention the polynucleotide comprises a
30 region encoding ribH polypeptides comprising a sequence set out in Table 1 [SEQ ID NO:1] which includes a full length gene, or a variant thereof.

In another particularly preferred embodiment of the invention there is a novel ribH protein from *Staphylococcus aureus* comprising the amino acid sequence of Table 1 [SEQ ID NO:2], or a variant thereof.

5 In accordance with another aspect of the invention there is provided an isolated nucleic acid molecule encoding a mature polypeptide expressible by the as *Staphylococcus aureus* WCUH29 strain, which nucleic acid is contained in the deposited strain.

A further aspect of the invention there are provided isolated nucleic acid molecules encoding ribH, particularly *Staphylococcus aureus* ribH, including mRNAs, cDNAs, genomic DNAs. Further embodiments of the invention include biologically, diagnostically, prophylactically, clinically
10 or therapeutically useful variants thereof, and compositions comprising the same.

In accordance with another aspect of the invention, there is provided the use of a polynucleotide of the invention for therapeutic or prophylactic purposes, in particular genetic immunization. Among the particularly preferred embodiments of the invention are naturally occurring allelic variants of ribH and polypeptides encoded thereby.

15 Another aspect of the invention there are provided novel polypeptides of *Staphylococcus aureus* referred to herein as ribH as well as biologically, diagnostically, prophylactically, clinically or therapeutically useful variants thereof, and compositions comprising the same.

Among the particularly preferred embodiments of the invention are variants of ribH polypeptide encoded by naturally occurring alleles of the ribH gene.

20 In a preferred embodiment of the invention there are provided methods for producing the aforementioned ribH polypeptides.

In accordance with yet another aspect of the invention, there are provided inhibitors to such polypeptides, useful as antibacterial agents, including, for example, antibodies.

25 In accordance with certain preferred embodiments of the invention, there are provided products, compositions and methods for assessing ribH expression, treating disease, assaying genetic variation, and administering a ribH polypeptide or polynucleotide to an organism to raise an immunological response against a bacteria, especially a *Staphylococcus aureus* bacteria.

30 In accordance with certain preferred embodiments of this and other aspects of the invention there are provided polynucleotides that hybridize to ribH polynucleotide sequences, particularly under stringent conditions.

In certain preferred embodiments of the invention there are provided antibodies against ribH polypeptides.

In other embodiments of the invention there are provided methods for identifying compounds which bind to or otherwise interact with and inhibit or activate an activity of a polypeptide or polynucleotide of the invention comprising: contacting a polypeptide or polynucleotide of the invention with a compound to be screened under conditions to permit binding to or other
 5 interaction between the compound and the polypeptide or polynucleotide to assess the binding to or other interaction with the compound, such binding or interaction being associated with a second component capable of providing a detectable signal in response to the binding or interaction of the polypeptide or polynucleotide with the compound; and determining whether the compound binds to or otherwise interacts with and activates or inhibits an activity of the polypeptide or polynucleotide
 10 by detecting the presence or absence of a signal generated from the binding or interaction of the compound with the polypeptide or polynucleotide.

In accordance with yet another aspect of the invention, there are provided ribH agonists and antagonists, preferably bacteriostatic or bacteriocidal agonists and antagonists.

In a further aspect of the invention there are provided compositions comprising a ribH
 15 polynucleotide or a ribH polypeptide for administration to a cell or to a multicellular organism.

Various changes and modifications within the spirit and scope of the disclosed invention will become readily apparent to those skilled in the art from reading the following descriptions and from reading the other parts of the present disclosure.

20 DESCRIPTION OF THE INVENTION

The invention relates to novel ribH polypeptides and polynucleotides as described in greater detail below. In particular, the invention relates to polypeptides and polynucleotides of a novel ribH of *Staphylococcus aureus*, which is related by amino acid sequence homology to *Bacillus amyloliquefaciens* ribH polypeptide. The invention relates especially to ribH having the nucleotide
 25 and amino acid sequences set out in Table 1 as SEQ ID NO: 1 and SEQ ID NO: 2 respectively ribH.

TABLE 1

ribH Polynucleotide and Polypeptide Sequences

30 (A) Sequences from *Staphylococcus aureus* ribH polynucleotide sequence [SEQ ID NO:1].
 5' -
 ATGAATTTTGAAGGTAAATTAATTGAAAAAGATTTGAAAGTTGCAATCGTAGTTAGTCGATTTAATGAT

TTTATCACTGGAAGATTACTTGAAGGTGCAAAAGATACTTTGATTGACATGATGTTAATGAAGACAAT
 ATTGATGTAGCATTTGTTCTGGTGCGTTTGAAATTCCTTTAGTAGCTAAAAAATTAGCCTCATCAGGA
 AATTATGATGCAATAATTACATTAGGATGCGTAATTCGCGGTGCTACGTCTCATTATGATTATGTTTGT
 AATGAAGTCGCGAAAGGTGTTTCTAAAGTAAATGATCAAATAATGTACCAGTCATATTTGGTATTTTA
 5 ACGACTGAAAGTATTGAACAAGCTGTGGAAAGAGCAGGTACGAAAGCTGGTAATAAAGGTGCCGAAGCA
 GCAGTAAGTGCAATTGAAATGGCTAATTTATTAAAATCTATAAAAGCATAG-3'

(B) *Staphylococcus aureus* ribH polypeptide sequence deduced from the polynucleotide sequence in this table [SEQ ID NO:2].

NH₂-

10 MNFEGKLI GKDLKVAIVVSRFNDFITGRLLGAKDTLIRHDVNEDNIDVAFVPGAFEIPLVAKKLASSG
 NYDAIITLGCVIRGATSHYDYVCNEVAKGVSKVNDQTNVPVIFGILTTESIEQAVERAGTKAGNKGAEA
 AVSAIEMANLLKSIKA-COOH

Deposited materials

15 A deposit containing a *Staphylococcus aureus* WCUH 29 strain has been deposited with the National Collections of Industrial and Marine Bacteria Ltd. (herein "NCIMB"), 23 St. Machar Drive, Aberdeen AB2 1RY, Scotland on 11 September 1995 and assigned NCIMB Deposit No. 40771, and referred to as *Staphylococcus aureus* WCUH29 on deposit. The *Staphylococcus aureus* strain deposit is referred to herein as "the deposited strain" or as "the DNA of the deposited strain."

20 The deposited strain contains the full length ribH gene. The sequence of the polynucleotides contained in the deposited strain, as well as the amino acid sequence of the polypeptide encoded thereby, are controlling in the event of any conflict with any description of sequences herein.

The deposit of the deposited strain has been made under the terms of the Budapest Treaty on the International Recognition of the Deposit of Micro-organisms for Purposes of Patent
 25 Procedure. The strain will be irrevocably and without restriction or condition released to the public upon the issuance of a patent. The deposited strain is provided merely as convenience to those of skill in the art and is not an admission that a deposit is required for enablement, such as that required under 35 U.S.C. §112.

A license may be required to make, use or sell the deposited strain, and compounds derived
 30 therefrom, and no such license is hereby granted.


One aspect of the invention there is provided an isolated nucleic acid molecule encoding a mature polypeptide expressible by the *Staphylococcus aureus* WCUH 29 strain contained in the deposited strain. Further provided by the invention are ribH nucleotide sequences of the DNA in the deposited strain and amino acid sequences encoded thereby. Also provided by the invention are ribH

polypeptide sequences isolated from the deposited strain and amino acid sequences derived therefrom.

Polypeptides

The polypeptides of the invention include a polypeptide of Table 1 [SEQ ID NO:2] (in particular the mature polypeptide) as well as polypeptides and fragments, particularly those which have the biological activity of ribH, and also those which have at least 70% identity to a polypeptide of Table 1 [SEQ ID NO:1] or the relevant portion, preferably at least 80% identity to a polypeptide of Table 1 [SEQ ID NO:2] and more preferably at least 90% similarity (more preferably at least 90% identity) to a polypeptide of Table 1 [SEQ ID NO:2] and still more preferably at least 95% similarity (still more preferably at least 95% identity) to a polypeptide of Table 1 [SEQ ID NO:2] and also include portions of such polypeptides with such portion of the polypeptide generally containing at least 30 amino acids and more preferably at least 50 amino acids.

The invention also includes polypeptides of the formula:



wherein, at the amino terminus, X is hydrogen, and at the carboxyl terminus, Y is hydrogen or a metal, R_1 and R_3 are any amino acid residue, m is an integer between 1 and 1000 or zero, n is an integer between 1 and 1000 or zero, and R_2 is an amino acid sequence of the invention, particularly an amino acid sequence selected from Table 1. In the formula above R_2 is oriented so that its amino terminal residue is at the left, bound to R_1 , and its carboxy terminal residue is at the right, bound to R_3 . Any stretch of amino acid residues denoted by either R group, where m and/or n is greater than 1, may be either a heteropolymer or a homopolymer, preferably a heteropolymer.

A fragment is a variant polypeptide having an amino acid sequence that entirely is the same as part but not all of the amino acid sequence of the aforementioned polypeptides. As with ribH polypeptides fragments may be "free-standing," or comprised within a larger polypeptide of which they form a part or region, most preferably as a single continuous region, a single larger polypeptide.

Preferred fragments include, for example, truncation polypeptides having a portion of an amino acid sequence of Table 1 [SEQ ID NO:2], or of variants thereof, such as a continuous series of residues that includes the amino terminus, or a continuous series of residues that includes the carboxyl terminus. Degradation forms of the polypeptides of the invention in a host cell, particularly a *Staphylococcus aureus*, are also preferred. Further preferred are fragments characterized by structural or functional attributes such as fragments that comprise alpha-helix and alpha-helix forming regions, beta-sheet and beta-sheet-forming regions, turn and turn-forming regions, coil and

coil-forming regions, hydrophilic regions, hydrophobic regions, alpha amphipathic regions, beta amphipathic regions, flexible regions, surface-forming regions, substrate binding region, and high antigenic index regions.

Also preferred are biologically active fragments which are those fragments that mediate activities of ribH, including those with a similar activity or an improved activity, or with a decreased undesirable activity. Also included are those fragments that are antigenic or immunogenic in an animal, especially in a human. Particularly preferred are fragments comprising receptors or domains of enzymes that confer a function essential for viability of *Staphylococcus aureus* or the ability to initiate, or maintain cause disease in an individual, particularly a human.

10 Variants that are fragments of the polypeptides of the invention may be employed for producing the corresponding full-length polypeptide by peptide synthesis; therefore, these variants may be employed as intermediates for producing the full-length polypeptides of the invention.

In addition to the standard single and triple letter representations for amino acids, the term "X" or "Xaa" may also be used in describing certain polypeptides of the invention. "X" and "Xaa" mean that any of the twenty naturally occurring amino acids may appear at such a designated position in the polypeptide sequence.

Polynucleotides

Another aspect of the invention relates to isolated polynucleotides, including the full length gene, that encode the ribH polypeptide having a deduced amino acid sequence of Table 1 [SEQ ID NO:2] and polynucleotides closely related thereto and variants thereof.

Using the information provided herein, such as a polynucleotide sequence set out in Table 1 [SEQ ID NO:1], a polynucleotide of the invention encoding ribH polypeptide may be obtained using standard cloning and screening methods, such as those for cloning and sequencing chromosomal DNA fragments from bacteria using *Staphylococcus aureus* WCUH 29 cells as starting material, followed by obtaining a full length clone. For example, to obtain a polynucleotide sequence of the invention, such as a sequence given in Table 1 [SEQ ID NO:1], typically a library of clones of chromosomal DNA of *Staphylococcus aureus* WCUH 29 in *E.coli* or some other suitable host is probed with a radiolabeled oligonucleotide, preferably a 17-mer or longer, derived from a partial sequence. Clones carrying DNA identical to that of the probe can then be distinguished using stringent conditions. By sequencing the individual clones thus identified with sequencing primers designed from the original sequence it is then possible to extend the sequence in both directions to determine the full gene sequence. Conveniently, such sequencing is performed

using denatured double stranded DNA prepared from a plasmid clone. Suitable techniques are described by Maniatis, T., Fritsch, E.F. and Sambrook et al., *MOLECULAR CLONING, A LABORATORY MANUAL*, 2nd Ed.; Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York (1989). (see in particular Screening By Hybridization 1.90 and Sequencing
5 Denatured Double-Stranded DNA Templates 13.70). Illustrative of the invention, the polynucleotide set out in Table 1 [SEQ ID NO:1] was discovered in a DNA library derived from *Staphylococcus aureus* WCUH 29.

The DNA sequence set out in Table 1 [SEQ ID NO:1] contains an open reading frame encoding a protein having about the number of amino acid residues set forth in Table 1 [SEQ ID
10 NO:2] with a deduced molecular weight that can be calculated using amino acid residue molecular weight values well known in the art. The polynucleotide of SEQ ID NO: 1, between nucleotide number 1 and the stop codon which begins at nucleotide number 463 of SEQ ID NO:1, encodes the polypeptide of SEQ ID NO:2.

RibH of the invention is structurally related to other proteins of the lumazine synthase (β -
15 subunit) family.

The invention provides a polynucleotide sequence identical over its entire length to a coding sequence in Table 1 [SEQ ID NO:1]. Also provided by the invention is the coding sequence for the mature polypeptide or a fragment thereof, by itself as well as the coding sequence for the mature polypeptide or a fragment in reading frame with other coding sequence, such as those encoding a
20 leader or secretory sequence, a pre-, or pro- or prepro- protein sequence. The polynucleotide may also contain non-coding sequences, including for example, but not limited to non-coding 5' and 3' sequences, such as the transcribed, non-translated sequences, termination signals, ribosome binding sites, sequences that stabilize mRNA, introns, polyadenylation signals, and additional coding sequence which encode additional amino acids. For example, a marker sequence that facilitates
25 purification of the fused polypeptide can be encoded. In certain embodiments of the invention, the marker sequence is a hexa-histidine peptide, as provided in the pQE vector (Qiagen, Inc.) and described in Gentz *et al.*, *Proc. Natl. Acad. Sci., USA* 86: 821-824 (1989), or an HA tag (Wilson *et al.*, *Cell* 37: 767 (1984). Polynucleotides of the invention also include, but are not limited to, polynucleotides comprising a structural gene and its naturally associated sequences that control gene
30 expression.

A preferred embodiment of the invention is a polynucleotide of comprising nucleotide 1 to the nucleotide immediately upstream of or including nucleotide 463 set forth in SEQ ID NO:1 of Table 1, both of which encode the ribH polypeptide.

The invention also includes polynucleotides of the formula:



wherein, at the 5' end of the molecule, X is hydrogen, and at the 3' end of the molecule, Y is hydrogen or a metal, R_1 and R_3 is any nucleic acid residue, m is an integer between 1 and 3000 or zero, n is an integer between 1 and 3000 or zero, and R_2 is a nucleic acid sequence of the invention, particularly a nucleic acid sequence selected from Table 1. In the polynucleotide formula above R_2 is oriented so that its 5' end residue is at the left, bound to R_1 , and its 3' end residue is at the right, bound to R_3 . Any stretch of nucleic acid residues denoted by either R group, where m and/or n is greater than 1, may be either a heteropolymer or a homopolymer, preferably a heteropolymer. In a preferred embodiment m and/or n is an integer between 1 and 1000.

It is most preferred that the polynucleotides of the inventions are derived from *Staphylococcus aureus*, however, they may preferably be obtained from organisms of the same taxonomic genus. They may also be obtained, for example, from organisms of the same taxonomic family or order.

The term "polynucleotide encoding a polypeptide" as used herein encompasses polynucleotides that include a sequence encoding a polypeptide of the invention, particularly a bacterial polypeptide and more particularly a polypeptide of the *Staphylococcus aureus* ribH having an amino acid sequence set out in Table 1 [SEQ ID NO:2]. The term also encompasses polynucleotides that include a single continuous region or discontinuous regions encoding the polypeptide (for example, interrupted by integrated phage or an insertion sequence or editing) together with additional regions, that also may contain coding and/or non-coding sequences.

The invention further relates to variants of the polynucleotides described herein that encode for variants of the polypeptide having a deduced amino acid sequence of Table 1 [SEQ ID NO:2]. Variants that are fragments of the polynucleotides of the invention may be used to synthesize full-length polynucleotides of the invention.

Further particularly preferred embodiments are polynucleotides encoding ribH variants, that have the amino acid sequence of ribH polypeptide of Table 1 [SEQ ID NO:2] in which several, a few, 5 to 10, 1 to 5, 1 to 3, 2, 1 or no amino acid residues are substituted, deleted or added, in any

combination. Especially preferred among these are silent substitutions, additions and deletions, that do not alter the properties and activities of ribH.

Further preferred embodiments of the invention are polynucleotides that are at least 70% identical over their entire length to a polynucleotide encoding ribH polypeptide having an amino acid sequence set out in Table 1 [SEQ ID NO:2], and polynucleotides that are complementary to such polynucleotides. Alternatively, most highly preferred are polynucleotides that comprise a region that is at least 80% identical over its entire length to a polynucleotide encoding ribH polypeptide and polynucleotides complementary thereto. In this regard, polynucleotides at least 90% identical over their entire length to the same are particularly preferred, and among these particularly preferred polynucleotides, those with at least 95% are especially preferred. Furthermore, those with at least 97% are highly preferred among those with at least 95%, and among these those with at least 98% and at least 99% are particularly highly preferred, with at least 99% being the more preferred.

Preferred embodiments are polynucleotides that encode polypeptides that retain substantially the same biological function or activity as the mature polypeptide encoded by a DNA of Table 1 [SEQ ID NO:1].

The invention further relates to polynucleotides that hybridize to the herein above-described sequences. In this regard, the invention especially relates to polynucleotides that hybridize under stringent conditions to the herein above-described polynucleotides. As herein used, the terms "stringent conditions" and "stringent hybridization conditions" mean hybridization will occur only if there is at least 95% and preferably at least 97% identity between the sequences. An example of stringent hybridization conditions is overnight incubation at 42°C in a solution comprising: 50% formamide, 5x SSC (150mM NaCl, 15mM trisodium citrate), 50 mM sodium phosphate (pH7.6), 5x Denhardt's solution, 10% dextran sulfate, and 20 micrograms/ml denatured, sheared salmon sperm DNA, followed by washing the hybridization support in 0.1x SSC at about 65°C. Hybridization and wash conditions are well known and exemplified in Sambrook, *et al.*, Molecular Cloning: A Laboratory Manual, Second Edition, Cold Spring Harbor, N.Y., (1989), particularly Chapter 11 therein.

The invention also provides a polynucleotide consisting essentially of a polynucleotide sequence obtainable by screening an appropriate library containing the complete gene for a polynucleotide sequence set forth in SEQ ID NO:1 under stringent hybridization conditions with a probe having the sequence of said polynucleotide sequence set forth in SEQ ID NO:1 or

a fragment thereof; and isolating said DNA sequence. Fragments useful for obtaining such a polynucleotide include, for example, probes and primers described elsewhere herein.

As discussed additionally herein regarding polynucleotide assays of the invention, for instance, polynucleotides of the invention as discussed above, may be used as a hybridization probe
5 for RNA, cDNA and genomic DNA to isolate full-length cDNAs and genomic clones encoding ribH and to isolate cDNA and genomic clones of other genes that have a high sequence similarity to the ribH gene. Such probes generally will comprise at least 15 bases. Preferably, such probes will have at least 30 bases and may have at least 50 bases. Particularly preferred probes will have at least 30 bases and will have 50 bases or less.

10 For example, the coding region of the ribH gene may be isolated by screening using a DNA sequence provided in Table 1 [SEQ ID NO: 1] to synthesize an oligonucleotide probe. A labeled oligonucleotide having a sequence complementary to that of a gene of the invention is then used to screen a library of cDNA, genomic DNA or mRNA to determine which members of the library the probe hybridizes to.

15 The polynucleotides and polypeptides of the invention may be employed, for example, as research reagents and materials for discovery of treatments of and diagnostics for disease, particularly human disease, as further discussed herein relating to polynucleotide assays.

Polynucleotides of the invention that are oligonucleotides derived from the sequences of Table 1 [SEQ ID NOS:1 or 2] may be used in the processes herein as described, but preferably
20 for PCR, to determine whether or not the polynucleotides identified herein in whole or in part are transcribed in bacteria in infected tissue. It is recognized that such sequences will also have utility in diagnosis of the stage of infection and type of infection the pathogen has attained.

The invention also provides polynucleotides that may encode a polypeptide that is the mature protein plus additional amino or carboxyl-terminal amino acids, or amino acids interior to the
25 mature polypeptide (when the mature form has more than one polypeptide chain, for instance). Such sequences may play a role in processing of a protein from precursor to a mature form, may allow protein transport, may lengthen or shorten protein half-life or may facilitate manipulation of a protein for assay or production, among other things. As generally is the case *in vivo*, the additional amino acids may be processed away from the mature protein by cellular enzymes.

30 A precursor protein, having the mature form of the polypeptide fused to one or more prosequences may be an inactive form of the polypeptide. When prosequences are removed such

inactive precursors generally are activated. Some or all of the prosequences may be removed before activation. Generally, such precursors are called proproteins.

In addition to the standard A, G, C, T/U representations for nucleic acid bases, the term "N" may also be used in describing certain polynucleotides of the invention. "N" means
5 that any of the four DNA or RNA bases may appear at such a designated position in the DNA or RNA sequence, except it is preferred that N is not a base that when taken in combination with adjacent nucleotide positions, when read in the correct reading frame, would have the effect of generating a premature termination codon in such reading frame.

In sum, a polynucleotide of the invention may encode a mature protein, a mature protein
10 plus a leader sequence (which may be referred to as a preprotein), a precursor of a mature protein having one or more prosequences that are not the leader sequences of a preprotein, or a preproprotein, which is a precursor to a proprotein, having a leader sequence and one or more prosequences, which generally are removed during processing steps that produce active and mature forms of the polypeptide.

15 Vectors, host cells, expression

The invention also relates to vectors that comprise a polynucleotide or polynucleotides of the invention, host cells that are genetically engineered with vectors of the invention and the production of polypeptides of the invention by recombinant techniques. Cell-free translation systems can also be employed to produce such proteins using RNAs derived from the DNA constructs of the invention.

20 For recombinant production, host cells can be genetically engineered to incorporate expression systems or portions thereof or polynucleotides of the invention. Introduction of a polynucleotide into the host cell can be effected by methods described in many standard laboratory manuals, such as Davis et al., *BASIC METHODS IN MOLECULAR BIOLOGY*, (1986) and Sambrook et al., *MOLECULAR CLONING: A LABORATORY MANUAL*, 2nd Ed., Cold Spring
25 Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989), such as, calcium phosphate transfection, DEAE-dextran mediated transfection, transvection, microinjection, cationic lipid-mediated transfection, electroporation, transduction, scrape loading, ballistic introduction and infection.

Representative examples of appropriate hosts include bacterial cells, such as streptococci,
30 staphylococci, enterococci *E. coli*, streptomyces and *Bacillus subtilis* cells; fungal cells, such as yeast cells and *Aspergillus* cells; insect cells such as *Drosophila* S2 and *Spodoptera* Sf9 cells;

animal cells such as CHO, COS, HeLa, C127, 3T3, BHK, 293 and Bowes melanoma cells; and plant cells.

A great variety of expression systems can be used to produce the polypeptides of the invention. Such vectors include, among others, chromosomal, episomal and virus-derived vectors, e.g., vectors derived from bacterial plasmids, from bacteriophage, from transposons, from yeast episomes, from insertion elements, from yeast chromosomal elements, from viruses such as baculoviruses, papova viruses, such as SV40, vaccinia viruses, adenoviruses, fowl pox viruses, pseudorabies viruses and retroviruses, and vectors derived from combinations thereof, such as those derived from plasmid and bacteriophage genetic elements, such as cosmids and phagemids. The expression system constructs may contain control regions that regulate as well as engender expression. Generally, any system or vector suitable to maintain, propagate or express polynucleotides and/or to express a polypeptide in a host may be used for expression in this regard. The appropriate DNA sequence may be inserted into the expression system by any of a variety of well-known and routine techniques, such as, for example, those set forth in Sambrook *et al.*, *MOLECULAR CLONING, A LABORATORY MANUAL*, (*supra*).

For secretion of the translated protein into the lumen of the endoplasmic reticulum, into the periplasmic space or into the extracellular environment, appropriate secretion signals may be incorporated into the expressed polypeptide. These signals may be endogenous to the polypeptide or they may be heterologous signals.

Polypeptides of the invention can be recovered and purified from recombinant cell cultures by well-known methods including ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography, and lectin chromatography. Most preferably, high performance liquid chromatography is employed for purification. Well known techniques for refolding protein may be employed to regenerate active conformation when the polypeptide is denatured during isolation and or purification.

Diagnostic Assays

This invention is also related to the use of the ribH polynucleotides of the invention for use as diagnostic reagents. Detection of ribH in a eukaryote, particularly a mammal, and especially a human, will provide a diagnostic method for diagnosis of a disease. Eukaryotes (herein also "individual(s)"), particularly mammals, and especially humans, particularly those infected or

suspected to be infected with an organism comprising the ribH gene may be detected at the nucleic acid level by a variety of techniques.

Nucleic acids for diagnosis may be obtained from an infected individual's cells and tissues, such as bone, blood, muscle, cartilage, and skin. Genomic DNA may be used directly for detection or may be amplified enzymatically by using PCR or other amplification technique prior to analysis. RNA, cDNA and genomic DNA may also be used in the same ways. Using amplification, characterization of the species and strain of prokaryote present in an individual, may be made by an analysis of the genotype of the prokaryote gene. Deletions and insertions can be detected by a change in size of the amplified product in comparison to the genotype of a reference sequence. Point mutations can be identified by hybridizing amplified DNA to labeled ribH polynucleotide sequences. Perfectly matched sequences can be distinguished from mismatched duplexes by RNase digestion or by differences in melting temperatures. DNA sequence differences may also be detected by alterations in the electrophoretic mobility of the DNA fragments in gels, with or without denaturing agents, or by direct DNA sequencing. See, e.g., Myers et al., *Science*, 230: 1242 (1985). Sequence changes at specific locations also may be revealed by nuclease protection assays, such as RNase and S1 protection or a chemical cleavage method. See, e.g., Cotton et al., *Proc. Natl. Acad. Sci., USA*, 85: 4397-4401 (1985).


Cells carrying mutations or polymorphisms in the gene of the invention may also be detected at the DNA level by a variety of techniques, to allow for serotyping, for example. For example, RT-PCR can be used to detect mutations. It is particularly preferred to used RT-PCR in conjunction with automated detection systems, such as, for example, GeneScan. RNA, cDNA or genomic DNA may also be used for the same purpose, PCR or RT-PCR. As an example, PCR primers complementary to a nucleic acid encoding ribH can be used to identify and analyze mutations. Examples of representative primers are shown below in Table 2.

Table 2

Primers for amplification of ribH polynucleotides

<u>SEQ ID NO</u>	<u>PRIMER SEQUENCE</u>
3	5'-AAGTTGCAATCGTAGTTAGTCG-3'
4	5'-GCTTCGGCACCTTTATTACCA-3'

The invention also includes primers of the formula:



wherein, at the 5' end of the molecule, X is hydrogen, and at the 3' end of the molecule, Y is hydrogen or a metal, R_1 and R_3 is any nucleic acid residue, m is an integer between 1 and 20 or zero, n is an integer between 1 and 20 or zero, and R_2 is a primer sequence of the invention, particularly a primer sequence selected from Table 2. In the polynucleotide formula above R_2 is oriented so that its 5' end residue is at the left, bound to R_1 , and its 3' end residue is at the right, bound to R_3 . Any stretch of nucleic acid residues denoted by either R group, where m and/or n is greater than 1, may be either a heteropolymer or a homopolymer, preferably a heteropolymer being complementary to a region of a polynucleotide of Table 1. In a preferred embodiment m and/or n is an integer between 1 and 10.

The invention further provides these primers with 1, 2, 3 or 4 nucleotides removed from the 5' and/or the 3' end. These primers may be used for, among other things, amplifying ribH DNA isolated from a sample derived from an individual. The primers may be used to amplify the gene isolated from an infected individual such that the gene may then be subject to various techniques for elucidation of the DNA sequence. In this way, mutations in the DNA sequence may be detected and used to diagnose infection and to serotype and/or classify the infectious agent.

The invention further provides a process for diagnosing, disease, preferably bacterial infections, more preferably infections by *Staphylococcus aureus*, comprising determining from a sample derived from an individual a increased level of expression of polynucleotide having a sequence of Table 1 [SEQ ID NO: 1]. Increased or decreased expression of ribH polynucleotide can be measured using any one of the methods well known in the art for the quantitation of polynucleotides, such as, for example, amplification, PCR, RT-PCR, RNase protection, Northern blotting and other hybridization methods.

In addition, a diagnostic assay in accordance with the invention for detecting over-expression of ribH protein compared to normal control tissue samples may be used to detect the presence of an infection, for example. Assay techniques that can be used to determine levels of a ribH protein, in a sample derived from a host are well-known to those of skill in the art. Such assay methods include radioimmunoassays, competitive-binding assays, Western Blot analysis and ELISA assays.

Antibodies

The polypeptides of the invention or variants thereof, or cells expressing them can be used as an immunogen to produce antibodies immunospecific for such polypeptides. "Antibodies" as used

herein includes monoclonal and polyclonal antibodies, chimeric, single chain, simianized antibodies and humanized antibodies, as well as Fab fragments, including the products of an Fab immunoglobulin expression library.

Antibodies generated against the polypeptides of the invention can be obtained by
5 administering the polypeptides or epitope-bearing fragments, analogues or cells to an animal, preferably a nonhuman, using routine protocols. For preparation of monoclonal antibodies, any technique known in the art that provides antibodies produced by continuous cell line cultures can be used. Examples include various techniques, such as those in Kohler, G. and Milstein, C., *Nature* 256: 495-497 (1975); Kozbor *et al.*, *Immunology Today* 4: 72 (1983); Cole *et al.*, pg. 77-96 in
10 *MONOCLONAL ANTIBODIES AND CANCER THERAPY*, Alan R. Liss, Inc. (1985).

Techniques for the production of single chain antibodies (U.S. Patent No. 4,946,778) can be adapted to produce single chain antibodies to polypeptides of this invention. Also, transgenic mice, or other organisms such as other mammals, may be used to express humanized antibodies.

Alternatively phage display technology may be utilized to select antibody genes with
15 binding activities towards the polypeptide either from repertoires of PCR amplified v-genes of lymphocytes from humans screened for possessing anti-ribH or from naive libraries (McCafferty, J. *et al.*, (1990), *Nature* 348, 552-554; Marks, J. *et al.*, (1992) *Biotechnology* 10, 779-783). The affinity of these antibodies can also be improved by chain shuffling (Clackson, T. *et al.*, (1991) *Nature* 352, 624-628).

20 If two antigen binding domains are present each domain may be directed against a different epitope - termed 'bispecific' antibodies.

The above-described antibodies may be employed to isolate or to identify clones expressing the polypeptides to purify the polypeptides by affinity chromatography.

Thus, among others, antibodies against ribH- polypeptide may be employed to treat
25 infections, particularly bacterial infections.

Polypeptide variants include antigenically, epitopically or immunologically equivalent variants that form a particular aspect of this invention. The term "antigenically equivalent derivative" as used herein encompasses a polypeptide or its equivalent which will be specifically recognized by certain antibodies which, when raised to the protein or polypeptide
30 according to the invention, interfere with the immediate physical interaction between pathogen and mammalian host. The term "immunologically equivalent derivative" as used herein encompasses a peptide or its equivalent which when used in a suitable formulation to raise

antibodies in a vertebrate, the antibodies act to interfere with the immediate physical interaction between pathogen and mammalian host.

The polypeptide, such as an antigenically or immunologically equivalent derivative or a fusion protein thereof is used as an antigen to immunize a mouse or other animal such as a rat or chicken. The fusion protein may provide stability to the polypeptide. The antigen may be associated, for example by conjugation, with an immunogenic carrier protein for example bovine serum albumin (BSA) or keyhole limpet haemocyanin (KLH). Alternatively a multiple antigenic peptide comprising multiple copies of the protein or polypeptide, or an antigenically or immunologically equivalent polypeptide thereof may be sufficiently antigenic to improve immunogenicity so as to obviate the use of a carrier.

Preferably, the antibody or variant thereof is modified to make it less immunogenic in the individual. For example, if the individual is human the antibody may most preferably be "humanized"; where the complementarity determining region(s) of the hybridoma-derived antibody has been transplanted into a human monoclonal antibody, for example as described in Jones, P. et al. (1986), *Nature* 321, 522-525 or Tempest et al., (1991) *Biotechnology* 9, 266-273.

The use of a polynucleotide of the invention in genetic immunization will preferably employ a suitable delivery method such as direct injection of plasmid DNA into muscles (Wolff et al., *Hum Mol Genet* 1992, 1:363, Manthorpe et al., *Hum. Gene Ther.* 1993:4, 419), delivery of DNA complexed with specific protein carriers (Wu et al., *J Biol Chem.* 1989: 264,16985), coprecipitation of DNA with calcium phosphate (Benvenisty & Reshef, *PNAS USA*, 1986:83,9551), encapsulation of DNA in various forms of liposomes (Kaneda et al., *Science* 1989:243,375), particle bombardment (Tang et al., *Nature* 1992, 356:152, Eisenbraun et al., *DNA Cell Biol* 1993, 12:791) and *in vivo* infection using cloned retroviral vectors (Seeger et al., *PNAS USA* 1984:81,5849).

Antagonists and agonists - assays and molecules

Polypeptides of the invention may also be used to assess the binding of small molecule substrates and ligands in, for example, cells, cell-free preparations, chemical libraries, and natural product mixtures. These substrates and ligands may be natural substrates and ligands or may be structural or functional mimetics. See, e.g., Coligan et al., *Current Protocols in Immunology* 1(2): Chapter 5 (1991).

The invention also provides a method of screening compounds to identify those which enhance (agonist) or block (antagonist) the action of ribH polypeptides or polynucleotides, particularly those compounds that are bacteriostatic and/or bacteriocidal. The method of screening may involve high-throughput techniques. For example, to screen for agonists or antagonists, a synthetic reaction mix, a cellular compartment, such as a membrane, cell envelope or cell wall, or a preparation of any thereof, comprising ribH polypeptide and a labeled substrate or ligand of such polypeptide is incubated in the absence or the presence of a candidate molecule that may be a ribH agonist or antagonist. The ability of the candidate molecule to agonize or antagonize the ribH polypeptide is reflected in decreased binding of the labeled ligand or decreased production of product from such substrate. Molecules that bind gratuitously, *i.e.*, without inducing the effects of ribH polypeptide are most likely to be good antagonists. Molecules that bind well and increase the rate of product production from substrate are agonists. Detection of the rate or level of production of product from substrate may be enhanced by using a reporter system. Reporter systems that may be useful in this regard include but are not limited to colorimetric labeled substrate converted into product, a reporter gene that is responsive to changes in ribH polynucleotide or polypeptide activity, and binding assays known in the art.

Another example of an assay for ribH antagonists is a competitive assay that combines ribH and a potential antagonist with ribH-binding molecules, recombinant ribH binding molecules, natural substrates or ligands, or substrate or ligand mimetics, under appropriate conditions for a competitive inhibition assay. RibH can be labeled, such as by radioactivity or a colorimetric compound, such that the number of ribH molecules bound to a binding molecule or converted to product can be determined accurately to assess the effectiveness of the potential antagonist.

Potential antagonists include small organic molecules, peptides, polypeptides and antibodies that bind to a polynucleotide or polypeptide of the invention and thereby inhibit or extinguish its activity. Potential antagonists also may be small organic molecules, a peptide, a polypeptide such as a closely related protein or antibody that binds the same sites on a binding molecule, such as a binding molecule, without inducing ribH-induced activities, thereby preventing the action of ribH by excluding ribH from binding.

Potential antagonists include a small molecule that binds to and occupies the binding site of the polypeptide thereby preventing binding to cellular binding molecules, such that normal biological activity is prevented. Examples of small molecules include but are not limited to small organic molecules, peptides or peptide-like molecules. Other potential antagonists include antisense

molecules (see Okano, *J. Neurochem.* 56: 560 (1991); *OLIGODEOXYNUCLEOTIDES AS ANTISENSE INHIBITORS OF GENE EXPRESSION*, CRC Press, Boca Raton, FL (1988), for a description of these molecules). Preferred potential antagonists include compounds related to and variants of ribH.

5 Each of the DNA sequences provided herein may be used in the discovery and development of antibacterial compounds. The encoded protein, upon expression, can be used as a target for the screening of antibacterial drugs. Additionally, the DNA sequences encoding the amino terminal regions of the encoded protein or Shine-Delgarno or other translation facilitating sequences of the respective mRNA can be used to construct antisense sequences to
10 control the expression of the coding sequence of interest.

The invention also provides the use of the polypeptide, polynucleotide or inhibitor of the invention to interfere with the initial physical interaction between a pathogen and mammalian host responsible for sequelae of infection. In particular the molecules of the invention may be used: in the prevention of adhesion of bacteria, in particular gram positive
15 bacteria, to mammalian extracellular matrix proteins on in-dwelling devices or to extracellular matrix proteins in wounds; to block ribH protein-mediated mammalian cell invasion by, for example, initiating phosphorylation of mammalian tyrosine kinases (Rosenshine *et al.*, *Infect. Immun.* 60:2211 (1992); to block bacterial adhesion between mammalian extracellular matrix proteins and bacterial ribH proteins that mediate tissue damage and; to block the normal
20 progression of pathogenesis in infections initiated other than by the implantation of in-dwelling devices or by other surgical techniques.

The antagonists and agonists of the invention may be employed, for instance, to inhibit and treat diseases.

Helicobacter pylori (herein *H. pylori*) bacteria infect the stomachs of over one-third of
25 the world's population causing stomach cancer, ulcers, and gastritis (International Agency for Research on Cancer (1994) Schistosomes, Liver Flukes and Helicobacter Pylori (International Agency for Research on Cancer, Lyon, France; <http://www.uicc.ch/ecp/ecp2904.htm>). Moreover, the international Agency for Research on Cancer recently recognized a cause-and-effect relationship between *H. pylori* and gastric adenocarcinoma, classifying the bacterium as
30 a Group I (definite) carcinogen. Preferred antimicrobial compounds of the invention (agonists and antagonists of ribH) found using screens provided by the invention, particularly broad-spectrum antibiotics, should be useful in the treatment of *H. pylori* infection. Such treatment

should decrease the advent of *H. pylori*-induced cancers, such as gastrointestinal carcinoma. Such treatment should also cure gastric ulcers and gastritis.

Vaccines

Another aspect of the invention relates to a method for inducing an immunological response in an individual, particularly a mammal which comprises inoculating the individual with ribH, or a fragment or variant thereof, adequate to produce antibody and/ or T cell immune response to protect said individual from infection, particularly bacterial infection and most particularly *Staphylococcus aureus* infection. Also provided are methods whereby such immunological response slows bacterial replication. Yet another aspect of the invention relates to a method of inducing immunological response in an individual which comprises delivering to such individual a nucleic acid vector to direct expression of ribH, or a fragment or a variant thereof, for expressing ribH, or a fragment or a variant thereof *in vivo* in order to induce an immunological response, such as, to produce antibody and/ or T cell immune response, including, for example, cytokine-producing T cells or cytotoxic T cells, to protect said individual from disease, whether that disease is already established within the individual or not. One way of administering the gene is by accelerating it into the desired cells as a coating on particles or otherwise. Such nucleic acid vector may comprise DNA, RNA, a modified nucleic acid, or a DNA/RNA hybrid.

A further aspect of the invention relates to an immunological composition which, when introduced into an individual capable or having induced within it an immunological response, induces an immunological response in such individual to a ribH or protein coded therefrom, wherein the composition comprises a recombinant ribH or protein coded therefrom comprising DNA which codes for and expresses an antigen of said ribH or protein coded therefrom. The immunological response may be used therapeutically or prophylactically and may take the form of antibody immunity or cellular immunity such as that arising from CTL or CD4+ T cells.

A ribH polypeptide or a fragment thereof may be fused with co-protein which may not by itself produce antibodies, but is capable of stabilizing the first protein and producing a fused protein which will have immunogenic and protective properties. Thus fused recombinant protein, preferably further comprises an antigenic co-protein, such as lipoprotein D from *Hemophilus influenzae*, Glutathione-S-transferase (GST) or beta-galactosidase, relatively large co-proteins which solubilize the protein and facilitate production and purification thereof. Moreover, the co-protein may act as an adjuvant in the sense of providing a generalized

stimulation of the immune system. The co-protein may be attached to either the amino or carboxy terminus of the first protein.

Provided by this invention are compositions, particularly vaccine compositions, and methods comprising the polypeptides or polynucleotides of the invention and
5 immunostimulatory DNA sequences, such as those described in Sato, Y. *et al.* Science 273: 352 (1996).

Also, provided by this invention are methods using the described polynucleotide or particular fragments thereof which have been shown to encode non-variable regions of bacterial cell surface proteins in DNA constructs used in such genetic immunization
10 experiments in animal models of infection with *Staphylococcus aureus* will be particularly useful for identifying protein epitopes able to provoke a prophylactic or therapeutic immune response. It is believed that this approach will allow for the subsequent preparation of monoclonal antibodies of particular value from the requisite organ of the animal successfully resisting or clearing infection for the development of prophylactic agents or therapeutic
15 treatments of bacterial infection, particularly *Staphylococcus aureus* infection, in mammals, particularly humans.

The polypeptide may be used as an antigen for vaccination of a host to produce specific antibodies which protect against invasion of bacteria, for example by blocking adherence of bacteria to damaged tissue. Examples of tissue damage include wounds in skin or
20 connective tissue caused, e.g., by mechanical, chemical or thermal damage or by implantation of indwelling devices, or wounds in the mucous membranes, such as the mouth, mammary glands, urethra or vagina.

The invention also includes a vaccine formulation which comprises an immunogenic recombinant protein of the invention together with a suitable carrier. Since the protein may be
25 broken down in the stomach, it is preferably administered parenterally, including, for example, administration that is subcutaneous, intramuscular, intravenous, or intradermal. Formulations suitable for parenteral administration include aqueous and non-aqueous sterile injection solutions which may contain anti-oxidants, buffers, bacteriostats and solutes which render the formulation isotonic with the bodily fluid, preferably the blood, of the individual; and aqueous
30 and non-aqueous sterile suspensions which may include suspending agents or thickening agents. The formulations may be presented in unit-dose or multi-dose containers, for example, sealed ampules and vials and may be stored in a freeze-dried condition requiring only the

addition of the sterile liquid carrier immediately prior to use. The vaccine formulation may also include adjuvant systems for enhancing the immunogenicity of the formulation, such as oil-in water systems and other systems known in the art. The dosage will depend on the specific activity of the vaccine and can be readily determined by routine experimentation.

5 While the invention has been described with reference to certain ribH protein, it is to be understood that this covers fragments of the naturally occurring protein and similar proteins with additions, deletions or substitutions which do not substantially affect the immunogenic properties of the recombinant protein.

Compositions, kits and administration

10 The invention also relates to compositions comprising the polynucleotide or the polypeptides discussed above or their agonists or antagonists. The polypeptides of the invention may be employed in combination with a non-sterile or sterile carrier or carriers for use with cells, tissues or organisms, such as a pharmaceutical carrier suitable for administration to a subject. Such compositions comprise, for instance, a media additive or a therapeutically effective amount of a polypeptide of the
15 invention and a pharmaceutically acceptable carrier or excipient. Such carriers may include, but are not limited to, saline, buffered saline, dextrose, water, glycerol, ethanol and combinations thereof. The formulation should suit the mode of administration. The invention further relates to diagnostic and pharmaceutical packs and kits comprising one or more containers filled with one or more of the ingredients of the aforementioned compositions of the invention.

20 Polypeptides and other compounds of the invention may be employed alone or in conjunction with other compounds, such as therapeutic compounds.

 The pharmaceutical compositions may be administered in any effective, convenient manner including, for instance, administration by topical, oral, anal, vaginal, intravenous, intraperitoneal, intramuscular, subcutaneous, intranasal or intradermal routes among others.

25 In therapy or as a prophylactic, the active agent may be administered to an individual as an injectable composition, for example as a sterile aqueous dispersion, preferably isotonic.

 Alternatively the composition may be formulated for topical application
for example in the form of ointments, creams, lotions, eye ointments, eye drops, ear drops, mouthwash, impregnated dressings and sutures and aerosols, and may contain appropriate
30 conventional additives, including, for example, preservatives, solvents to assist drug penetration, and emollients in ointments and creams. Such topical formulations may also contain compatible conventional carriers, for example cream or ointment bases, and ethanol or

oleyl alcohol for lotions. Such carriers may constitute from about 1% to about 98% by weight of the formulation; more usually they will constitute up to about 80% by weight of the formulation.

For administration to mammals, and particularly humans, it is expected that the daily
5 dosage level of the active agent will be from 0.01 mg/kg to 10 mg/kg, typically around 1 mg/kg. The physician in any event will determine the actual dosage which will be most suitable for an individual and will vary with the age, weight and response of the particular individual. The above dosages are exemplary of the average case. There can, of course, be individual instances where higher or lower dosage ranges are merited, and such are within the scope of
10 this invention.

In-dwelling devices include surgical implants, prosthetic devices and catheters, i.e., devices that are introduced to the body of an individual and remain in position for an extended time. Such devices include, for example, artificial joints, heart valves, pacemakers, vascular grafts, vascular catheters, cerebrospinal fluid shunts, urinary catheters, continuous ambulatory
15 peritoneal dialysis (CAPD) catheters.

The composition of the invention may be administered by injection to achieve a systemic effect against relevant bacteria shortly before insertion of an in-dwelling device. Treatment may be continued after surgery during the in-body time of the device. In addition, the composition could also be used to broaden perioperative cover for any surgical technique to
20 prevent bacterial wound infections, especially *Staphylococcus aureus* wound infections.

Many orthopaedic surgeons consider that humans with prosthetic joints should be considered for antibiotic prophylaxis before dental treatment that could produce a bacteremia. Late deep infection is a serious complication sometimes leading to loss of the prosthetic joint and is accompanied by significant morbidity and mortality. It may therefore be possible to
25 extend the use of the active agent as a replacement for prophylactic antibiotics in this situation.

In addition to the therapy described above, the compositions of this invention may be used generally as a wound treatment agent to prevent adhesion of bacteria to matrix proteins exposed in wound tissue and for prophylactic use in dental treatment as an alternative to, or in conjunction with, antibiotic prophylaxis.

30 Alternatively, the composition of the invention may be used to bathe an indwelling device immediately before insertion. The active agent will preferably be present at a concentration of 1 µg/ml to 10mg/ml for bathing of wounds or indwelling devices.

A vaccine composition is conveniently in injectable form. Conventional adjuvants may be employed to enhance the immune response. A suitable unit dose for vaccination is 0.5-5 microgram/kg of antigen, and such dose is preferably administered 1-3 times and with an interval of 1-3 weeks. With the indicated dose range, no adverse toxicological effects will be observed with the compounds of the invention which would preclude their administration to suitable individuals.

Each reference disclosed herein is incorporated by reference herein in its entirety. Any patent application to which this application claims priority is also incorporated by reference herein in its entirety.

10 GLOSSARY

The following definitions are provided to facilitate understanding of certain terms used frequently herein.

"Disease(s)" means and disease caused by or related to infection by a bacteria, including disease, such as, infections of the upper respiratory tract (e.g., otitis media, bacterial tracheitis, acute epiglottitis, thyroiditis), lower respiratory (e.g., empyema, lung abscess), cardiac (e.g., infective endocarditis), gastrointestinal (e.g., secretory diarrhoea, splenic abscess, retroperitoneal abscess), CNS (e.g., cerebral abscess), eye (e.g., blepharitis, conjunctivitis, keratitis, endophthalmitis, preseptal and orbital cellulitis, dacryocystitis), kidney and urinary tract (e.g., epididymitis, intrarenal and perinephric abscess, toxic shock syndrome), skin (e.g., impetigo, folliculitis, cutaneous abscesses, cellulitis, wound infection, bacterial myositis) bone and joint (e.g., septic arthritis, osteomyelitis).

"Host cell" is a cell which has been transformed or transfected, or is capable of transformation or transfection by an exogenous polynucleotide sequence.

"Identity," as known in the art, is a relationship between two or more polypeptide sequences or two or more polynucleotide sequences, as determined by comparing the sequences. In the art, "identity" also means the degree of sequence relatedness between polypeptide or polynucleotide sequences, as the case may be, as determined by the match between strings of such sequences. "Identity" and "similarity" can be readily calculated by known methods, including but not limited to those described in (*Computational Molecular Biology*, Lesk, A.M., ed., Oxford University Press, New York, 1988; *Biocomputing: Informatics and Genome Projects*, Smith, D.W., ed., Academic Press, New York, 1993; *Computer Analysis of Sequence Data*, Part I, Griffin, A.M., and Griffin, H.G., eds., Humana Press, New Jersey, 1994; *Sequence Analysis in Molecular Biology*, von Heinje, G., Academic Press, 1987; and *Sequence Analysis Primer*,

Gribskov, M. and Devereux, J., eds., M Stockton Press, New York, 1991; and Carillo, H., and Lipman, D., *SIAM J. Applied Math.*, 48: 1073 (1988). Preferred methods to determine identity are designed to give the largest match between the sequences tested. Methods to determine identity and similarity are codified in publicly available computer programs.

5 Preferred computer program methods to determine identity and similarity between two sequences include, but are not limited to, the GCG program package (Devereux, J., et al., *Nucleic Acids Research* 12(1): 387 (1984)), BLASTP, BLASTN, and FASTA (Atschul, S.F. et al., *J. Molec. Biol.* 215: 403-410 (1990)). The BLAST X program is publicly available from NCBI and other sources (*BLAST Manual*, Altschul, S., et al., NCBI NLM NIH

10 Bethesda, MD 20894; Altschul, S., et al., *J. Mol. Biol.* 215: 403-410 (1990)). As an illustration, by a polynucleotide having a nucleotide sequence having at least, for example, 95% "identity" to a reference nucleotide sequence of SEQ ID NO: 1 it is intended that the nucleotide sequence of the polynucleotide is identical to the reference sequence except that the polynucleotide sequence may include up to five point mutations per each 100 nucleotides of the

15 reference nucleotide sequence of SEQ ID NO: 1. In other words, to obtain a polynucleotide having a nucleotide sequence at least 95% identical to a reference nucleotide sequence, up to 5% of the nucleotides in the reference sequence may be deleted or substituted with another nucleotide, or a number of nucleotides up to 5% of the total nucleotides in the reference sequence may be inserted into the reference sequence. These mutations of the reference

20 sequence may occur at the 5' or 3' terminal positions of the reference nucleotide sequence or anywhere between those terminal positions, interspersed either individually among nucleotides in the reference sequence or in one or more contiguous groups within the reference sequence. Analogously, by a polypeptide having an amino acid sequence having at least, for example, 95% identity to a reference amino acid sequence of SEQ ID NO:2 is intended that the amino

25 acid sequence of the polypeptide is identical to the reference sequence except that the polypeptide sequence may include up to five amino acid alterations per each 100 amino acids of the reference amino acid of SEQ ID NO: 2. In other words, to obtain a polypeptide having an amino acid sequence at least 95% identical to a reference amino acid sequence, up to 5% of the amino acid residues in the reference sequence may be deleted or substituted with another

30 amino acid, or a number of amino acids up to 5% of the total amino acid residues in the reference sequence may be inserted into the reference sequence. These alterations of the reference sequence may occur at the amino or carboxy terminal positions of the reference

amino acid sequence or anywhere between those terminal positions, interspersed either individually among residues in the reference sequence or in one or more contiguous groups within the reference sequence.

"Isolated" means altered "by the hand of man" from its natural state, *i.e.*, if it occurs in nature, it has been changed or removed from its original environment, or both. For example, a polynucleotide or a polypeptide naturally present in a living organism is not "isolated," but the same polynucleotide or polypeptide separated from the coexisting materials of its natural state is "isolated", as the term is employed herein. Moreover, a polynucleotide or polypeptide that is introduced into an organism by transformation, genetic manipulation or by any other recombinant method is "isolated" even if it is still present in said organism, which organism may be living or non-living.

"Polynucleotide(s)" generally refers to any polyribonucleotide or polydeoxribonucleotide, which may be unmodified RNA or DNA or modified RNA or DNA. "Polynucleotide(s)" include, without limitation, single- and double-stranded DNA, DNA that is a mixture of single- and double-stranded regions or single-, double- and triple-stranded regions, single- and double-stranded RNA, and RNA that is mixture of single- and double-stranded regions, hybrid molecules comprising DNA and RNA that may be single-stranded or, more typically, double-stranded, or triple-stranded regions, or a mixture of single- and double-stranded regions. In addition, "polynucleotide" as used herein refers to triple-stranded regions comprising RNA or DNA or both RNA and DNA. The strands in such regions may be from the same molecule or from different molecules. The regions may include all of one or more of the molecules, but more typically involve only a region of some of the molecules. One of the molecules of a triple-helical region often is an oligonucleotide. As used herein, the term "polynucleotide(s)" also includes DNAs or RNAs as described above that contain one or more modified bases. Thus, DNAs or RNAs with backbones modified for stability or for other reasons are "polynucleotide(s)" as that term is intended herein. Moreover, DNAs or RNAs comprising unusual bases, such as inosine, or modified bases, such as tritylated bases, to name just two examples, are polynucleotides as the term is used herein. It will be appreciated that a great variety of modifications have been made to DNA and RNA that serve many useful purposes known to those of skill in the art. The term "polynucleotide(s)" as it is employed herein embraces such chemically, enzymatically or metabolically modified forms of polynucleotides, as well as the chemical forms of DNA and RNA characteristic of viruses and cells, including, for example, simple

and complex cells. "Polynucleotide(s)" also embraces short polynucleotides often referred to as oligonucleotide(s).

"Polypeptide(s)" refers to any peptide or protein comprising two or more amino acids joined to each other by peptide bonds or modified peptide bonds. "Polypeptide(s)" refers to both short chains, commonly referred to as peptides, oligopeptides and oligomers and to longer chains generally referred to as proteins. Polypeptides may contain amino acids other than the 20 gene encoded amino acids. "Polypeptide(s)" include those modified either by natural processes, such as processing and other post-translational modifications, but also by chemical modification techniques. Such modifications are well described in basic texts and in more detailed monographs, as well as in a voluminous research literature, and they are well known to those of skill in the art. It will be appreciated that the same type of modification may be present in the same or varying degree at several sites in a given polypeptide. Also, a given polypeptide may contain many types of modifications. Modifications can occur anywhere in a polypeptide, including the peptide backbone, the amino acid side-chains, and the amino or carboxyl termini. Modifications include, for example, acetylation, acylation, ADP-ribosylation, amidation, covalent attachment of flavin, covalent attachment of a heme moiety, covalent attachment of a nucleotide or nucleotide derivative, covalent attachment of a lipid or lipid derivative, covalent attachment of phosphatidylinositol, cross-linking, cyclization, disulfide bond formation, demethylation, formation of covalent cross-links, formation of cysteine, formation of pyroglutamate, formylation, gamma-carboxylation, glycosylation, GPI anchor formation, hydroxylation, iodination, methylation, myristoylation, oxidation, proteolytic processing, phosphorylation, prenylation, racemization, glycosylation, lipid attachment, sulfation, gamma-carboxylation of glutamic acid residues, hydroxylation and ADP-ribosylation, selenoylation, sulfation, transfer-RNA mediated addition of amino acids to proteins, such as arginylation, and ubiquitination. See, for instance, *PROTEINS - STRUCTURE AND MOLECULAR PROPERTIES*, 2nd Ed., T. E. Creighton, W. H. Freeman and Company, New York (1993) and Wold, F., Posttranslational Protein Modifications: Perspectives and Prospects, pgs. 1-12 in *POSTTRANSLATIONAL COVALENT MODIFICATION OF PROTEINS*, B. C. Johnson, Ed., Academic Press, New York (1983); Seifter et al., *Meth. Enzymol.* 182:626-646 (1990) and Rattan et al., *Protein Synthesis: Posttranslational Modifications and Aging*, Ann. N.Y. Acad. Sci. 663: 48-62 (1992). Polypeptides may be branched or cyclic, with or without branching. Cyclic, branched and branched circular polypeptides may result from post-translational natural processes and may be made by entirely synthetic methods, as well.

“Variant(s)” as the term is used herein, is a polynucleotide or polypeptide that differs from a reference polynucleotide or polypeptide respectively, but retains essential properties. A typical variant of a polynucleotide differs in nucleotide sequence from another, reference polynucleotide. Changes in the nucleotide sequence of the variant may or may not alter the amino acid sequence of a polypeptide encoded by the reference polynucleotide. Nucleotide changes may result in amino acid substitutions, additions, deletions, fusions and truncations in the polypeptide encoded by the reference sequence, as discussed below. A typical variant of a polypeptide differs in amino acid sequence from another, reference polypeptide. Generally, differences are limited so that the sequences of the reference polypeptide and the variant are closely similar overall and, in many regions, identical. A variant and reference polypeptide may differ in amino acid sequence by one or more substitutions, additions, deletions in any combination. A substituted or inserted amino acid residue may or may not be one encoded by the genetic code. A variant of a polynucleotide or polypeptide may be a naturally occurring such as an allelic variant, or it may be a variant that is not known to occur naturally. Non-naturally occurring variants of polynucleotides and polypeptides may be made by mutagenesis techniques, by direct synthesis, and by other recombinant methods known to skilled artisans.

EXAMPLES

The examples below are carried out using standard techniques, which are well known and routine to those of skill in the art, except where otherwise described in detail. The examples are illustrative, but do not limit the invention.

Example 1 Strain selection, Library Production and Sequencing

The polynucleotide having a DNA sequence given in Table 1 [SEQ ID NO:1] was obtained from a library of clones of chromosomal DNA of *Staphylococcus aureus* in *E. coli*. The sequencing data from two or more clones containing overlapping *Staphylococcus aureus* DNAs was used to construct the contiguous DNA sequence in SEQ ID NO:1. Libraries may be prepared by routine methods, for example:

Methods 1 and 2 below.

Total cellular DNA is isolated from *Staphylococcus aureus* WCUH 29 according to standard procedures and size-fractionated by either of two methods.

30 Method 1

Total cellular DNA is mechanically sheared by passage through a needle in order to size-fractionate according to standard procedures. DNA fragments of up to 11kbp in size are

rendered blunt by treatment with exonuclease and DNA polymerase, and EcoRI linkers added. Fragments are ligated into the vector Lambda ZapII that has been cut with EcoRI, the library packaged by standard procedures and *E.coli* infected with the packaged library. The library is amplified by standard procedures.

5 **Method 2**

Total cellular DNA is partially hydrolyzed with a one or a combination of restriction enzymes appropriate to generate a series of fragments for cloning into library vectors (e.g., RsaI, PstI, AluI, BshI235I), and such fragments are size-fractionated according to standard procedures. EcoRI linkers are ligated to the DNA and the fragments then ligated into the
10 vector Lambda ZapII that have been cut with EcoRI, the library packaged by standard procedures, and *E.coli* infected with the packaged library. The library is amplified by standard procedures.

Example 2 ribH Characterization

15 **Characterization of *Staphylococcus aureus* ribH Gene Expression**
Gene expression *in vivo*

Recently several novel approaches have been described which purport to follow global gene expression during infection (Chuang, S. et al., (1993); Mahan, M.J. et al., *Science* 259:686-688 (1993); Hensel, M. et al., *Science* 269:400-403 (1995). These new techniques
20 have so far been demonstrated with gram negative pathogen infections but not with infections with gram positives presumably because the much slower development of global transposon mutagenesis and suitable vectors needed for these strategies in these organisms, and in the case of that process described by Chuang, S. et al., *J. Bacteriol.* 175:2026-2036 (1993) the difficulty of isolating suitable quantities of bacterial RNA free of mammalian RNA derived
25 from the infected tissue to furnish bacterial RNA labelled to sufficiently high specific activity.

The present invention employs a novel technology to determine gene expression in the pathogen at different stages of infection of the mammalian host.

Use of the technology of the present invention enables identification of bacterial genes transcribed during infection, inhibitors of which would have utility in anti-bacterial therapy.
30 Specific inhibitors of such gene transcription or of the subsequent translation of the resultant mRNA or of the function of the corresponding expressed proteins would have utility in anti-bacterial therapy.

The determination of expression during infection of a gene from *Staphylococcus aureus* WCUH29

Necrotic fatty tissue from a four day groin infection or kidney from a seven day pyelonephritis infection of *Staphylococcus aureus* WCUH29 in the mouse is efficiently
5 disrupted and processed in the presence of acid phenol and detergent to provide a mixture of animal and bacterial RNA. By freezing the tissue immediately in liquid nitrogen, and processing the tissue samples while still frozen, changes in the population of bacterial mRNA is minimized. The resultant total RNA is free of DNA and protein (including RNAases and DNAases). The optimal conditions for disruption and processing to give high yields of
10 bacterial mRNA with transcripts of long length are followed by reverse transcribing the resulting mRNA to cDNA and amplified with ORF-specific primers for a bacterial gene known to be expressed constitutively and at low copy number in *Staphylococcus aureus* WCUH29. Aspects of this example II, part b, are modifications of a published protocol (Cheung, et al.; Anal Biochem (1994) 222:511-514).

15

a) Isolation of tissue infected with *Staphylococcus aureus* WCUH29 from murine models of infection

i) Isolation of tissue infected with *Staphylococcus aureus* WCUH29 from a murine thigh lesion model of infection. 10 ml. volumes of sterile nutrient broth (No.2 Oxoid)
20 are seeded with isolated, individual colonies of *Staphylococcus aureus* WCUH29 from an agar culture plate. The cultures are incubated aerobically (static culture) at 37°C for 16-20 hours. 4 week old mice (female, 18g-22g, strain MF1) are each infected by subcutaneous injection of 0.5ml. of this broth culture of *Staphylococcus aureus* WCUH29 (diluted in broth to approximately 10^8 cfu/ml.) into the anterior, right lower quadrant (groin area). Mice should be
25 monitored regularly during the first 24 hours after infection, then daily until termination of study. Animals with signs of systemic infection, i.e. lethargy, ruffled appearance, isolation from group, should be monitored closely and if signs progress to moribundancy, the animal should be culled immediately.

Visible external signs of lesion development will be seen 24-48h after infection.
30 Examination of the abdomen of the animal will show the raised outline of the abscess beneath the skin. The localised lesion should remain in the right lower quadrant, but may occasionally spread to the left lower quadrant, and superiorly to the thorax. On occasions, the abscess may rupture through the overlying skin layers. In such cases the affected animal should be culled immediately and the tissues sampled if possible. Failure to cull the animal may result in the
35 necrotic skin tissue overlying the abscess being sloughed off, exposing the abdominal muscle wall.

Approximately 96 hours after infection, animals are killed using carbon dioxide asphyxiation. To minimise delay between death and tissue processing /storage, mice should be killed individually rather than in groups. The dead animal is placed onto its back and the fur swabbed liberally with 70% alcohol. An initial incision using scissors is made through the skin of the abdominal left lower quadrant, travelling superiorly up to, then across the thorax. The incision is completed by cutting inferiorly to the abdominal lower right quadrant. Care should be taken not to penetrate the abdominal wall. Holding the skin flap with forceps, the skin is gently pulled away from the abdomen. The exposed abscess, which covers the peritoneal wall but generally does not penetrate the muscle sheet completely, is excised, taking care not to puncture the viscera

The abscess/muscle sheet and other infected tissue may require cutting in sections, prior to flash-freezing in liquid nitrogen, thereby allowing easier storage in plastic collecting vials.

ii) **Isolation of tissue infected with *Staphylococcus aureus* WCUH29 from a murine model of hematogenous pyelonephritis.** Overnight cultures of *S. aureus* WCUH29 are started from single colonies in 5 ml of tryptic soy broth (TSB) and grown at 37C with shaking. The cultures are then washed twice in sterile phosphate-buffered saline (PBS) and diluted to an A600 = 0.3. Male CD-1 mice (18 - 20g) are infected with 0.2 ml of this suspension by tail vein inoculation using a 30g needle attached to a tuberculin syringe. Each mouse receives approximately 4×10^7 bacteria in this fashion. Mice are monitored daily for signs of illness, and usually within 48 hours show signs of lethargy, ruffled fur, sluggishness; animals which appear moribund are euthanized prior to the end of the experiment.

All animals are euthanized via carbon dioxide overdose seven days post-infection. The animal is placed on its back and swabbed with ethanol, and then with RNAZap, and instruments are swabbed as well. The abdominal cavity is opened and the kidneys aseptically removed, cut into four pieces, and placed in cryovials which are immediately frozen in liquid nitrogen.

b) Isolation of *Staphylococcus aureus* WCUH29 RNA from infected tissue samples

Infected tissue samples, in 2-ml cryo-storage tubes, are removed from liquid nitrogen storage for immediate processing of the frozen tissue. In a microbiological safety cabinet the samples are disrupted up to eight at a time. To disrupt the bacteria within the tissue sample, 50-100 mg of the tissue is transferred to a FastRNA tube containing a silica/ceramic matrix (BIO101). Immediately, 1 ml of extraction reagents (FastRNA reagents, BIO101) are added to give a sample to reagent volume ratio of approximately 1 to 20. The tubes are shaken in a

reciprocating shaker (FastPrep FP120, BIO101) at a setting of 5.5 to 6 for 20-120 sec. The crude RNA preparation is extracted with chloroform/isoamyl alcohol, and precipitated with DEPC-treated/Isopropanol Precipitation Solution (BIO101). RNA preparations are stored in this isopropanol solution at -80°C if necessary. The RNA is pelleted (12,000g for 10 min.), washed with 75% ethanol (v/v in DEPC-treated water), air-dried for 5-10 min, and resuspended in 0.1 ml of DEPC-treated water.

Quality of the RNA isolated is assessed by the ability to detect bacterial transcripts up to 2 kb in length by RT-PCR (as described in section below). To demonstrate the isolation of bacterial RNA from the infected tissue, samples of RNA are reverse transcribed, and the presence of a constitutively expressed gene is detected through the use of quantitative PCR in the presence of a TaqMan probe (as described below).

c) The removal of DNA from *Staphylococcus aureus* WCUH29-derived RNA

DNA was removed from 50 microgram samples of RNA by a 30 minute treatment at 37°C with 10 units of RNAase-free DNAaseI (GeneHunter) in the buffer supplied in a final volume of 57 microliters.

The DNAase was inactivated and removed by phenol:chloroform extraction. RNA was precipitated with 5 microliters of 3 M NaOAc and 200 microliters 100% EtOH, and pelleted by centrifugation at 12,000g for 10 minutes. The RNA is pelleted (12,000g for 10 min.), washed with 75% ethanol (v/v in DEPC-treated water), air-dried for 5-10 min, and resuspended in 10-20 microliters of DEPC-treated water. RNA yield is quantitated by OD₂₆₀ after 1:1000 dilution of the cleaned RNA sample. RNA is stored at -80°C if necessary and reverse-transcribed within one week.

d) The preparation of cDNA from RNA samples derived from infected tissue

10 microliter samples of DNAase treated RNA are reverse transcribed using a SuperScript Preamplification System for First Strand cDNA Synthesis kit (Gibco BRL, Life Technologies) according to the manufacturers instructions. 1 nanogram of random hexamers is used to prime each reaction. Controls without the addition of SuperScriptII reverse transcriptase are also run. Both +/-RT samples are treated with RNaseH before proceeding to the PCR reaction .

e) The use of PCR to determine the quality of bacterial RNA derived from infected tissue

Long transcripts, which are expected to be of low copy number within the bacterial cell, such as penicillin-binding protein 2 (PBP2), are reverse transcribed with random primers as described above and amplified by the following PCR method using ORF-specific primers, in

order to ascertain the quality, represented by length amplified, of the mRNA obtained during extraction and purification.

5 PCR reactions are set up on ice in 0.2 ml tubes in a total volume of 50 ul by adding the following components [final concentration]: AmpliTaq PCR Buffer II (1X), 1.5mM MgCL₂, 1 mM dNTPs, 0.5uM forward primer, 0.5uM reverse primer, and 2 ul reverse-transcribed RNA. PCR reactions are run on a PE GeneAmp PCR System 9600 with an initial step of 94°C for 2min, followed by 35 cycles of 94 °C for 30sec, 42 °C for 30sec and 72 °C for 30sec, followed by a final extensison at 72 °C for 7min.

10

f) The use of PCR to determine the presence of a bacterial cDNA species

PCR reactions are set up as described above using 0.5 microM each of the ORF specific forward and reverse primers.

15 PCR product in 20 microliter aliquots are separated by electrophoresis through 1 to 1.5% 1 x TBE agarose gels or 10% 1x TBE acrylamide gels. PCR product is visualized by staining the gel with ethidium bromide. Size estimates are made by comparison to a 100 bp DNA Ladder (Gibco BRL, Life Technologies). Alternatively, if the PCR products are conveniently labelled by the use of a labelled PCR primer (e.g. labelled at the 5'end with a dye) a suitable aliquot of the PCR product is run out on a polyacrylamide sequencing gel and its
20 presence and quantity detected using a suitable gel scanning system (e.g. ABI Prism™ 377 Sequencer using GeneScan™ software as supplied by Perkin Elmer).

RT/PCR controls may include +/- reverse transcriptase reactions, 16s rRNA primers or DNA specific primer pairs designed to produce PCR products from non-transcribed *Staphylococcus aureus* WCUH29 genomic sequences.

25 To test the efficiency of the primer pairs they are used in DNA PCR with *Staphylococcus aureus* WCUH29 total DNA. PCR reactions are set up and run as described above using approx. 1 microgram of DNA in place of the cDNA and 35 cycles of PCR.

Primer pairs which fail to give the predicted sized product in either DNA PCR or RT/PCR are PCR failures and as such are uninformative. Of those which give the correct size
30 product with DNA PCR two classes are distinguished in RT/PCR: 1. Genes which are not transcribed *in vivo* reproducibly fail to give a product in RT/PCR; and 2. Genes which are transcribed *in vivo* reproducibly give the correct size product in RT/PCR and show a stronger signal in the +RT samples than the signal (if at all present) in -RT controls.

35 **g) The use of PCR and fluorogenic probes to determine the presence of a bacterial cDNA species**

Specific sequence detection occurs by amplification of target sequences in the PE Applied Biosystems 7700 Sequence Detection System in the presence of an oligonucleotide probe labeled at the 5' and 3' ends with a reporter and quencher fluorescent dye, respectively (FQ probe), which anneals between the two PCR primers. Only specific product will be
 5 detected when the probe is bound between the primers. As PCR amplification proceeds, the 5'-nuclease activity of Taq polymerase initially cleaves the reporter dye from the probe. The signal generated when the reporter dye is physically separated from the quencher dye is detected by measuring the signal with an attached CCD camera. Each signal generated equals one probe cleaved which corresponds to amplification of one target strand

10 PCR reactions are set up using the PE Applied Biosystem TaqMan PCR Core Reagent Kit according to the instructions supplied such that each reaction contains 5 microliters 10X PCR Buffer II, 7 microliters 25 mM MgCl₂, 5 microliters 300 nM forward primer, 5 microliters reverse primer, 5 microliters specific FQ probe, 1 microliter each 10 mM dATP, 10 mM dCTP, 10 mM dGTP and 20 mM dUTP, 13.25 microliters distilled water, 0.5 microliters
 15 AmpErase UNG, and 0.25 microliters AmpliTaq DNA polymerase to give a total volume of 45 microliters.

Amplification proceeds under the following thermal cycling conditions: 50°C hold for 2 minutes, 95°C hold for 10 minutes, 40 cycles of 95°C for 15 seconds and 60°C for 1 minute, followed by a 25°C hold until sample is retrieved. Detection occurs real-time. Data is
 20 collected at the end of the reaction.

Based on these analyses it was discovered that the *Staphylococcus aureus ribH* gene was transcribed *in vivo*.

25

Riboflavin Biosynthesis Operon

This ORF is part of an operon which encodes genes *ribG*, *ribB*, *ribA* and *ribH*. Gene *ribG* starts at nucleotide 1 and ends at nucleotide 1029. Gene *ribB* starts at nucleotide 1036 and
 30 ends at nucleotide 1668. Gene *ribA* starts at nucleotide 1679 and ends at nucleotide 2860. Gene *ribH* starts at nucleotide 2873 and ends at nucleotide 3337. The operon is as follows (SEQ ID NO: 5)

ATGGATTATGCGATTCAACTTGCAAATATGGTACAAGGTCAAACAGGTGTTAATCCACCCGTTGGCGCT
 35 GTTGTAGTTAATGAAGGTAGGATTGTTGGTATTGGTGCACACTTGAGAAAAGGTGACAAGCATGCGGAG
 GTTCAAGCACTTGATATGGCACAACAAAATGCTGAAGGTGCGACGATTTATATTACGTTAGAGCCATGT

AGTCATTTTGGTTCAACACCACCCTGTGTTAAACAAAATTATTGATTGTAAGATAGCAAAAGTAGTATAC
GCAACAAAAGACAATTCGTTAGACACACATGGTGATGAGACGTTACGGGCTCACGGTATTGAGGTTGAA
TGCGTTGATGATGAACGGGCATCACAATTATACCAAGACTTTTTTAAAGCAAAAGCAAAGCAACTGCCA
CAAATTACAGTGAAAGTATCTGCAAGTTTAGATGGTAAACAAGCGAATGATAATGGACAAAGTCAATGG
5 ATTACTAACAAAGAGGTTAAACAAGATGTCTATAAGTTAAGACATCGACACGACGCAGTGTTAACTGGA
AGACGTACAGTTGAATTAGATGATCCACAATATACTACACGTATTCAAGATGGAAAAACCCTATAAAA
GTAATATTGTCTAAGTCTGGGAATATTCATTTTAATCAGCAAATTTATCAAGATGAATCAACACCAATT
TGGATATATACTGAAAATCCAAATTTAAACAAGCAATCAAACACATATTGAAATTATTTACTTGAAGTCT
TGTGATTTAAACAACAATTCCTCACAATTTATATAAAAGAGGAGTTGGAACTTTGCTAGTCGAGGCAGGT
10 CCAACCACTACTTCAGAATTCCTCATCTATTATATAGATGAATTTATTCTCTATTATGCCCCGAAATTA
ATTGGCGGATCTGGAAATTATCAATTTTATCAAACAAATGATGTGATTGAGATACCAGATGCGAACC
TTTGAAATTGTTCAATTCGAGTTATTAAATCAAATGTTAAATTAACTTTACGAAAGAAGTGATGATGC
ATGTTTACTGGCATCGTTGAAGAAATAGGTGTCGTTAAAGTGTTCAAATTCGTCAATCAGTCAGGACG
ATTGAAATCGAAGCACATAAGATTACGGCAGATATGCATATTGGTGATTCCATCAGTGTTAATGGTGCA
15 TGTTTAACAGTGATTGATTTTGATCAGACATCTTTTACTGTTCAAGTAATTTAAAGCACTGAAAAATAAA
ACCTATTTAGCAGATGTTAAGCGACAATCAGAAGTAAATTTAGAGCGTGCCATGAGTGGAACGGTAGG
TTTGGTGGACATTTTGTCTCGGTCATGTAGATGAACTAGGAACAGTTTCAAAAATAAATGAAACAGCC
AATGCCAAAATTATTACGATTCAATGTAGCCAACATATTAATAATCAGTTAGTTAAGCAAGGGTCTATT
ACTGTGGATGGTGTAAGTCTAACGGTATTTGATAAGCATGATAACAGTTTGTGACATTCATCTTATTCCA
20 GAAACGAGGCGTTCAACGATTTTATCATCCAAAAAATTAGGAGATAAAGTACATTTAGAAACAGACGTT
TTGTTTAAATATGTTGAAAATATTTTAAATAAAGATAAAGACCAATTATCTGTAGATAAATTAAGAGCA
TTTGGTTTTTAGGAGGGGTAGCATGCAATTCGATAATATTGACAGTGCTTTAATGGCTTTAAAAAATGG
AGAAACAATTATTGTAGTAGATGATGAGAATCGTGAAAATGAAGGTGATTTAGTAGCGGTTACTGAATG
GATGAACGATAATACCATTAATTTTATGGCGAAAGAAGCAAGGGGATTAATATGCGCACCAGTGCTAA
25 AGATATTGCACAACGTTTGGATTTGGTACAAATGGTTGATGATAACTCCGACATCTTTGGTACGCAATT
TACAGTGAGTATTGATCATGTAGATACAACAACAGGAATTAGTGCTTATGAACGTACATTGACTGCCAA
AAAGCTCATTGATCCTAGTAGTGAAGCTAAAGATTTTAATCGTCCTGGTCATTTATTTCCATTAGTAGC
ACAAGATAAAGGCGTATTAGCTAGAAATGGACACACAGAAGCGGCTGTTGATTTAGCTAAACTTACTGG
TGCCAAGCCCGCTGGTGTCAATTTGTGAGATTATGAATGATGACGGCAGATGGCGAAAGGACAAGATTT
30 ACAAATTTTAAAGAAAAACATCAATTAAAGATGATTACGATTGATGATTTAATTGAATATCGTAAAAA
ATTAGAACCAGAAATTGAATTTAAGGCAAAAGTGAAAATGCCTACAGATTTTCGGAACATTTGATATGTA
TGGTTTTTAAAGCGACATACACAGATGAAGAGATAGTTGTACTGACAAAAGGTGCAATTCGACAACATGA
AAATGTACGCTTACATTCTGCGTGCCTTACAGGCGATATTTCCATAGTCAACGTTGTGATTGTGGTGC
TCAACTTGAATCGTCTATGAAGTATATCAATGAACATGGTGGCATGATTATTTATCTACCTCAAGAAGG
35 TCGTGGCATAGGATTGTTAAACAAATTACGCGCATATGAATTAATTGAGCAAGGATATGATACAGTAAC
TGCAAATTTAGCATTAGGTTTTGATGAAGATTTACGAGATTATCATATTGCTGCACAGATTTTAAATA

TTTTAACATCGAACATATCAATTTATTAAGTAATAATCCAAGTAAATTTGAGGGATTAAAACAATATGG
CATTGATATTGCAGAAAGAATTGAAGTTATCGTACCAGAAACGGTACATAATCATGATTATATGGTAAC
GAAAAAATAAAAAATGGGTCATTTAATATAGGAGGACTTTAACATGAATTTTGAAGGTAAATTAATTGG
AAAAGATTTGAAAGTTGCAATCGTAGTTAGTCGATTTAATGATTTTATCACTGGAAGATTACTTGAAGG
5 TGCAAAAGATACTTTGATTCGACATGATGTTAATGAAGACAATATTGATGTAGCATTTGTTCCCTGGTGC
GTTTGAAATTCCTTTAGTAGCTAAAAAATTAGCCTCATCAGGAAATTATGATGCAATAATTACATTAGG
ATGCGTAATTCGCGGTGCTACGTCTCATTATGATTATGTTTGTAATGAAGTGCGAAAGGTGTTTCTAAA
GTAAATGATCAAATAATGTACCAGTCATATTTGGTATTTTAACGACTGAAAGTATTGAACAAGCTGTG
GAAAGAGCAGGTACGAAAGCTGGTAATAAAGGTGCCGAAGCAGCAGTAAGTGCAATTGAAATGGCTAAT
10 TTATTAAAATCTATAAAAGCATAG

What is claimed is:

1. An isolated polynucleotide comprising a polynucleotide having at least a 70% identity to a polynucleotide encoding a polypeptide comprising the amino acid sequence of SEQ ID NO:2.
2. An isolated polynucleotide comprising a polynucleotide having at least a 70% identity to a polynucleotide encoding the same mature polypeptide expressed by the ribH gene contained in the *Staphylococcus aureus* of the deposited strain.
3. An isolated polynucleotide comprising a polynucleotide encoding a polypeptide comprising an amino acid sequence which is at least 70% identical to the amino acid sequence of SEQ ID NO:2.
4. An isolated polynucleotide that is complementary to the polynucleotide of claim 1
5. The polynucleotide of Claim 1 wherein the polynucleotide is DNA or RNA
6. The polynucleotide of Claim 1 comprising the nucleic acid sequence set forth in SEQ ID NO:1.
7. The polynucleotide of Claim 1 comprising nucleotide 1 to the stop codon which begins at nucleotide number 463 set forth in SEQ ID NO:1.
8. The polynucleotide of Claim 1 which encodes a polypeptide comprising the amino acid sequence of SEQ ID NO:2.
9. A vector comprising the polynucleotide of Claim 1.
10. A host cell comprising the vector of Claim 9.
11. A process for producing a polypeptide comprising: expressing from the host cell of Claim 10 a polypeptide encoded by said DNA.
12. A process for producing a ribH polypeptide or fragment comprising culturing a host of claim 10 under conditions sufficient for the production of said polypeptide or fragment.
13. A polypeptide comprising an amino acid sequence which is at least 70% identical to the amino acid sequence of SEQ ID NO:2.
14. A polypeptide comprising an amino acid sequence as set forth in SEQ ID NO:2.
15. An antibody against the polypeptide of claim 14.
16. An antagonist which inhibits the activity or expression of the polypeptide of claim 14.

17. A method for the treatment of an individual in need of ribH polypeptide comprising: administering to the individual a therapeutically effective amount of the polypeptide of claim 14.
18. A method for the treatment of an individual having need to inhibit ribH polypeptide comprising: administering to the individual a therapeutically effective amount of the antagonist of Claim 14.
19. A process for diagnosing a disease related to expression or activity of the polypeptide of claim 14 in an individual comprising:
- (a) determining a nucleic acid sequence encoding said polypeptide, and/or
 - (b) analyzing for the presence or amount of said polypeptide in a sample derived from the individual.
20. A method for identifying compounds which interact with and inhibit or activate an activity of the polypeptide of claim 14 comprising:
- contacting a composition comprising the polypeptide with the compound to be screened under conditions to permit interaction between the compound and the polypeptide to assess the interaction of a compound, such interaction being associated with a second component capable of providing a detectable signal in response to the interaction of the polypeptide with the compound;
 - and determining whether the compound interacts with and activates or inhibits an activity of the polypeptide by detecting the presence or absence of a signal generated from the interaction of the compound with the polypeptide.
21. A method for inducing an immunological response in a mammal which comprises inoculating the mammal with ribH polypeptide of claim 14, or a fragment or variant thereof, adequate to produce antibody and/or T cell immune response to protect said animal from disease.
22. A method of inducing immunological response in a mammal which comprises delivering a nucleic acid vector to direct expression of ribH polypeptide of claim 14, or fragment or a variant thereof, for expressing said ribH polypeptide, or a fragment or a variant thereof *in vivo* in order to induce an immunological response to produce antibody and/ or T cell immune response to protect said animal from disease.

SEQUENCE LISTING

5 <110> SmithKline Beecham Corporation
 <120> ribH
 <130> GM10123
 10 <150> 08/977,553
 <151> 1997-11-25
 <160> 5
 15 <170> FastSEQ for Windows Version 3.0
 <210> 1
 <211> 465
 20 <212> DNA
 <213> Staphylococcus aureus
 <400> 1
 25 atgaattttg aaggtaaatt aattggaaaa gatttgaaag ttgcaatcgt agttagtcga
 60
 tttaatgatt ttatcactgg aagattactt gaagggtgcaa aagatacttt gattcgacat
 120
 gatgttaatg aagacaatat tgatgtagca tttgttcctg gtgcgtttga aattccttta
 180
 30 gtagctaaaa aattagcctc atcaggaaat tatgatgcaa taattacatt aggatgcgta
 240
 attcgcggtg ctacgtctca ttatgattat gtttgtaatg aagtcgcgaa aggtgtttct
 300
 aaagtaaattg atcaaactaa tgtaccagtc atatttggtg ttttaacgac tgaaagtatt
 35 360
 gaacaagctg tggaaagagc aggtacgaaa gctggtaata aagggtgccga agcagcagta
 420
 agtgcaattg aaatggctaa tttattaaaa tctataaaag catag
 465
 40 <210> 2

<211> 154

<212> PRT

<213> Staphylococcus aureus

5

<400> 2

Met Asn Phe Glu Gly Lys Leu Ile Gly Lys Asp Leu Lys Val Ala Ile
 1 5 10 15
 Val Val Ser Arg Phe Asn Asp Phe Ile Thr Gly Arg Leu Leu Glu Gly
 20 25 30
 10 Ala Lys Asp Thr Leu Ile Arg His Asp Val Asn Glu Asp Asn Ile Asp
 35 40 45
 Val Ala Phe Val Pro Gly Ala Phe Glu Ile Pro Leu Val Ala Lys Lys
 50 55 60
 Leu Ala Ser Ser Gly Asn Tyr Asp Ala Ile Ile Thr Leu Gly Cys Val
 15 65 70 75 80
 Ile Arg Gly Ala Thr Ser His Tyr Asp Tyr Val Cys Asn Glu Val Ala
 85 90 95
 Lys Gly Val Ser Lys Val Asn Asp Gln Thr Asn Val Pro Val Ile Phe
 100 105 110
 20 Gly Ile Leu Thr Thr Glu Ser Ile Glu Gln Ala Val Glu Arg Ala Gly
 115 120 125
 Thr Lys Ala Gly Asn Lys Gly Ala Glu Ala Ala Val Ser Ala Ile Glu
 130 135 140
 Met Ala Asn Leu Leu Lys Ser Ile Lys Ala
 25 145 150

<210> 3

<211> 22

<212> DNA

30

<213> Staphylococcus aureus

<400> 3

aagttgcaat cgtagttagt cg

22

35

<210> 4

<211> 21

<212> DNA

<213> Staphylococcus aureus

40

<400> 4

gcttcggcac ctttattacc a

21

<210> 5
<211> 3336
<212> DNA
5 <213> Staphylococcus aureus

<400> 5
atggattatg cgattcaact tgcaaatatg gtacaaggtc aaacaggtgt taatccaccc
60
10 gttggcgctg ttgtagttaa tgaaggtagg attgttggtg ttggtgcaca cttgagaaaa
120
ggtgacaagc atgcggaggt tcaagcactt gatatggcac aacaaaatgc tgaaggtagc
180
acgatttata ttacgttaga gccatgtagt cattttgggt caacaccacc ctgtgttaac
15 240
aaaattattg attgtaagat agcaaaagta gtatacgcaa caaaagacaa ttcgttagac
300
acacatggtg atgagacggt acgggctcac ggtattgagg ttgaatgcgt tgatgatgaa
360
20 cgggcatcac aattatacca agactttttt aaagcaaaag caaagcaact gccacaaatt
420
acagtgaaag tatctgcaag tttagatggt aaacaagcga atgataatgg acaaagtcaa
480
tggtacta acaaagaggt taaacaagat gtctataagt taagacatcg acacgacgca
25 540
gtgttaactg gaagacgtac agttgaatta gatgatccac aatatactac acgtattcaa
600
gatggaaaaa accctataaa agtaatatg tctaagtctg ggaatattca ttttaatcag
660
30 caaatattatc aagatgaatc aacaccaatt tggatatata ctgaaaatcc aaatttaaca
720
agcaatcaaa cacatattga aattattttac ttgaagtctt gtgatttaac aacaattctt
780
cacaatttat ataaaagagg agttggaact ttgctagtcg aggcaggtcc aaccactact
35 840
tcagaattct ccatctatta tatagatgaa tttattctct attatgcccc gaaattaatt
900
ggcggatctg gaaattatca attttatcaa acaaatgatg tgattgagat accagatgcg
960
40 aaccaatttg aaattgttca ttccgagtta ttaaatcaaa atgttaaatt aactttacga
1020

aagaagtgat gatgcatggt tactggcatc gttgaagaaa taggtgtcgt taaaagtgtt
 1080
 caaattcgtc aatcagtcag gacgattgaa atcgaagcac ataagattac ggcagatatg
 1140
 5 catattggtg attccatcag tgttaatggt gcatgtttta cagtgattga ttttgatcag
 1200
 acatctttta ctgttcaagt aattaaaagc actgaaaata aaacctatct agcagatggt
 1260
 aagcgacaat cagaagtaaa tttagagcgt gccatgagtg gtaacggtag gtttggtgga
 10 1320
 cattttgtcc tcggtcatgt agatgaacta ggaacagttt caaaaataaa tgaaacagcc
 1380
 aatgccaaaa ttattacgat tcaatgtagc caacatatta ataatcagtt agttaagcaa
 1440
 15 gggcttatta ctgtggatgg tgtaagtcta acggtatttg ataagcatga taacagtttt
 1500
 gacattcatc ttattccaga aacgagggcgt tcaacgattt tatcatccaa aaaattagga
 1560
 gataaagtac atttagaaac agacgttttg tttaaatatg ttgaaaatat tttaaataaa
 20 1620
 gataaagacc aattatctgt agataaatta agagcatttg gtttttagga ggggtagcat
 1680
 gcaattcgat aatattgaca gtgctttaat ggctttaaaa aatggagaaa caattattgt
 1740
 25 agtagatgat gagaatcgtg aaaatgaagg tgatttagta gcggttactg aatggatgaa
 1800
 cgataatacc attaatctta tggcgaaaga agcaagggga ttaatatgcg caccagtgtc
 1860
 taaagatatt gcacaacggt tggatttggt acaaagggt gatgataact ccgacatctt
 30 1920
 tggtagcga tttacagtga gtattgatca ttagatata acaacaggaa ttagtgctta
 1980
 tgaacgtaca ttgactgcca aaaagctcat tgatcctagt agtgaagcta aagattttta
 2040
 35 tcgtcctggt catttatctc cattagtagc acaagataaa ggcgtattag ctagaaatgg
 2100
 acacacagaa gcggctgttg atttagctaa acttactggt gccagcccg ctggtgtcat
 2160
 ttgtgagatt atgaatgatg acggcacgat ggcgaaagga caagatttac aaaattttta
 40 2220

agaaaaacat caattaaaga tgattacgat tgatgattta attgaatatc gtaaaaaatt
2280
agaaccagaa attgaattta aggcaaaagt gaaaatgcct acagatttcg gaacatttga
2340
5 tatgtatggt tttaaagcga catacacaga tgaagagata gttgtactga caaaagggtgc
2400
aattcgacaa catgaaaatg tacgcttaca ttctgcgtgc cttacaggcg atattttcca
2460
tagtcaacgt tgtgattgtg gtgctcaact tgaatcgtct atgaagtata tcaatgaaca
10 2520
tggtggcatg attattttatc tacctcaaga aggtcgtggc ataggattgt taaacaaatt
2580
acgcgcatat gaattaattg agcaaggata tgatacagta actgcaaatt tagcattagg
2640
15 ttttgatgaa gatttacgag attatcatat tgctgcacag attttaaaat attttaacat
2700
cgaacatatc aattttattaa gtaataatcc aagtaaattt gagggattaa aacaatatgg
2760
cattgatatt gcagaaagaa ttgaagttat cgtaccagaa acggtacata atcatgatta
20 2820
tatggtaacg aaaaaaataa aaatgggtca tttaatatag gaggacttta acatgaattt
2880
tgaaggtaaa ttaattggaa aagatttgaa agttgcaatc gtagttagtc gatttaatga
2940
25 ttttatcact ggaagattac ttgaagggtgc aaaagatact ttgattcgac atgatgttaa
3000
tgaagacaat attgatgtag catttgttcc tgggtgcgttt gaaattcctt tagtagctaa
3060
aaaattagcc tcatcaggaa attatgatgc aataattaca ttaggatgcg taattcgcgg
30 3120
tgctacgtct cattatgatt atgtttgtaa tgaagtgcga aagggtgttc taaagtaa
3180
gatcaaacta atgtaccagt catatttgggt attttaacga ctgaaagtat tgaacaagct
3240
35 gtggaaagag caggtagcga agctggtaat aaagggtgccg aagcagcagt aagtgaatt
3300
gaaatggcta atttattaaa atctataaaa gcatag
3336

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/25096

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : C12P 21/00; C12N 1/20, 1/21, 5/12, 15/00

US CL : 536/23.1; 435/320.1, 71.1, 172.3, 240.27

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 536/23.1; 435/320.1, 71.1, 172.3, 240.27

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Please See Extra Sheet.

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	GUSAROV, I. I. ET AL. The genes for riboflavin biosynthesis in <i>Bacillus amyloliquefaciens</i> : Primary structure, organization, and regulation. <i>Molecular Biology</i> . 1997, Vol. 31, No. 3, pages 370-376, see entire document.	1-12
X	LEE, C. Y. ET AL. Riboflavin synthesis genes are linked with the lux operon of <i>Photobacterium phosphoreum</i> . <i>J. Bacteriology</i> . April 1994, Vol. 176, No. 7, pages 2100-2104, see entire document.	1-12

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*A* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

27 JANUARY 1999

Date of mailing of the international search report

09 FEB 1999

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

NITA M. MINNIFIELD

Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US98/25096

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

Please See Extra Sheet.

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:
1-12

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US98/25096

B. FIELDS SEARCHED

Electronic data bases consulted (Name of data base and where practicable terms used):

BIOSIS, EMBASE, MEDLINE, JAPIO, BIOTECHDS, WPIDS, CABA, APS, MSPARS SEQUENCE SEARCH

inventor names, ribH, polynucleotide, nucleic acid, isolat?, DNA, genomic DN, staphylococcus, 6, 7-dimethyl-8-ribityllumazine synthase, riboflavin

BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING

This ISA found multiple inventions as follows:

This application contains the following inventions or groups of inventions which are not so linked as to form a single inventive concept under PCT Rule 13.1. In order for all inventions to be searched, the appropriate additional search fees must be paid.

Group I, claims 1-12, drawn to polynucleotides.

Group II, claims 13-14, drawn to a polypeptide.

Group III, claim 15, drawn to an antibody.

Group IV, claim 16, drawn to an antagonist.

Group V, claim 17, drawn to a method of treatment comprising administering a polypeptide.

Group VI, claim 18, drawn to a method of treatment comprising administering an antagonist.

Group VII, claim 19a, drawn to a process for diagnosis comprising determining a nucleic acid sequence.

Group VIII, claim 19b, drawn to a process for diagnosis comprising analyzing a polypeptide.

Group IX, claim 20, drawn to a method for identifying compounds.

Group X, claim 21, drawn to a method for inducing an immunological response which comprises inoculation with polypeptide.

Group XI, claim 22, drawn to a method for inducing an immunological response which comprises inoculation with a nucleic acid vector.

The inventions listed as Groups I-XI do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons: the special technical feature of Group I is considered to be the polynucleotides; the special technical feature of Group II is considered to be the polypeptide; the special technical feature of Group III is considered to be the antibody; the special technical feature of

Group IV is considered to be the antagonist; the special technical feature of Group V is considered to be a method of treatment comprising administering a polypeptide; the special technical feature of Group VI is considered to be a method of treatment comprising administering an antagonist; the special technical feature of Group VII is considered to be a process for diagnosis comprising determining a nucleic acid sequence; the special technical feature of Group VIII is considered to be a process for diagnosis comprising analyzing a polypeptide; the special technical feature of Group IX is considered to be a method for identifying compounds; the special technical feature of Group X is considered to be a method for inducing an immunological response which comprises inoculation with polypeptide; the special technical feature of Group XI is considered to be a method for inducing an immunological response which comprises inoculation with a nucleic acid vector.

Accordingly, Groups I-XI are not so linked by the same or correspondingly special technical feature as to form a single general inventive concept.